Oregon Department of Education





Rick Sherman

Oregon Department of Education Farm to School / School Garden Coordinator rick.sherman@state.or.us For more resources visit: www.ode.state.or.us/go/f2sgardens

Oregon Department of Education School Garden Food Safety Program



Oregon Department of Education

School Garden Food Safety Program



Instructions:

Read the companion guide first. It will guide you step by step through this training and documentation system.

Table of Contents

Page 1:	School Garden Food Safety Program Companion Guide
Tab 1:	School Garden Startup Checklists
Tab 2:	School Garden Weekly Checklists
Tab 3:	USDA Best Practices - "Handling Fresh Produce in Schools"
Tab 4:	School Garden Staff Training & Log
Tab 5:	School Garden Student Training & Log
Tab 6:	Composting Logs

Startup Checklists

Weekly Checklists

Handling Fresh Produce

Staff Training /Log

Student Training /Log

Composting Logs

Oregon Department of Education

School Garden Food Safety Training & Documentation Manual

~Companion Guide~



In Oregon, many people are resistant to serve food in their cafeterias that have been obtained from school gardens. Sometimes there are misconceptions that serving this food might not be allowed, or that serving the food from the gardens is unsafe. This is not the case, as long as some simple food handling procedures are followed.

Fresh produce served in schools from school gardens in Oregon requires no formal food safety inspection for "minimally processed" fruits and vegetables. "Minimally processed" means *processing the food that will not fundamentally change the properties of the food*. Examples of minimally processed food include cutting, chopping and freezing. Other foods in a school garden setting served from sources such as raising chicken eggs, or processing beef or poultry at the school would require a lengthy inspection and certification process, and could not legally be served in schools without this inspection and certification.

All food served to the public or in school cafeterias from the school garden must be handled following the same food handling procedures that are required by the local County Health Department in Oregon. Other than local health regulations, there are no requirements or regulations for school gardens. However, it is important to know that EVERYONE associated with a school garden can be liable for the safety of the students and staff that work in the garden or consume the food. For this reason, the Oregon Department of Education has developed this garden safety program as a way to potentially minimize risks associated with serving food from school gardens.

This manual is written with Oregon laws in mind. In other states, there are differing laws and rules concerning the subjects discussed here. This manual is meant to be used as a tool to help minimize risks associated with school gardens, and is not intended to be the "final word" on all aspects of safe school gardening.

Note that these recommendations refer to food safety practices for both harvesting from the school garden for consumption by others (in the garden or the cafeteria) and harvesting directly for personal consumption. Safe food handling is always important, but is most critical when handling food consumed by others that is typically eaten raw (without a cooking step to kill pathogens).

The following is an overview of what's in the manual. Refer to the other sections to see detailed steps and explanations of the material

Keep in mind the following:

- Documentation is a standard practice in the restaurant and cafeteria worlds. There's a saying that goes: "If it's not documented, IT DIDN'T HAPPEN!" In other words, you have to prove it happened by documenting it.
- If there's ever an issue with a foodborne illness outbreak in your area, the school garden could be at risk if there isn't proper documentation available. If there are good records, it will help tremendously by showing that your garden is doing everything possible to minimize risks and ensure the safety of your garden.
- This system will only work if you actually DO what is documented.

CONTENTS of the Garden Safety Program: (all items can be found in the GARDEN SAFETY section of the ODE Farm to School/School Garden website) <u>www.ode.state.or.us/go/f2sgardens</u> -Click on "School Gardens" box

- SGFSP Companion Guide (this document)
- School Garden Startup Checklist (complete each Fall and Spring)
- School Garden WEEKLY CHECKLIST
- USDA "Best Practices Handling Fresh Produce in Schools"
- School Garden Staff Training & Log
- School Garden Student Training & Log
- Composting logs

STARTUP CHECKLIST (Tab 1):

- 1. This checklist is to be completed twice per year. It's recommended that the list be completed once in the Fall (by September 30), and in the Spring (by April 30). Complete a separate checklist each time. Enter the date in the appropriate space.
- 2. Enter the name of the designated adult that is in charge of the garden.
- 3. Enter at least one designated adult with current food handler certification that will be in charge of handling the produce from the garden. This is usually the Kitchen Manager or Cook, although it technically could be the same person. Oregon Food Handler permits can be obtained online at www.orfoodhandlers.com and are inexpensive. In a foodservice environment all food handlers need to obtain a food handler permit. It's conceded that it's impractical to have every student to have a food handler permit, but at the least the leader or leaders of the garden who are in charge of harvesting food should get one.
- 4. Identify soil history of the garden site. This will help determine areas of the garden that aren't suitable for growing food or may need special amendments.
- 5. Soil should be tested for lead, arsenic and other major metals and contaminants. If you are using the existing soil (rather than bringing in new soil), a one-time test should be sufficient. Test new soil as it is added to the garden. It's recommended to contact your local OSU county extension office for referral to people who test soil. You can find your local extension office/agent here: <u>http://extension.oregonstate.edu/find-us</u>. Test for lead and other contaminants that are identified through soil review. The extension

office should be able to assist you in determining what concerns there may be with your soil. Soil with content greater than or equal to 400 mg of lead per kg is considered potentially hazardous. Here are some other tips:

- Sample depth should be 4 to 6 inches, retrieve soil from several areas in your garden.
- Contact the local utility companies (or call 811 the national "call before you dig" number) a few days before digging to ensure that you avoid gas or electric lines.
- 6. Is municipal, potable (safe for drinking) water used for irrigation (watering)? If so, then the water is being monitored by the municipality and is certainly adequate for irrigation.
- 7. If not (if you are using well water, water collection systems, etc.), test the water at least once a year. This is not an expensive process. It is recommended that you contact your local OSU county extension office for referral to people who test water. Remember, water collection systems/containers should be cleaned and sanitized regularly.
- 8. The school garden plot should be positioned so that it is not in the path of runoff from agricultural areas (such as livestock or dairy farms, etc.), parking lots, roads, or other sources of potential contamination.
- 9. ODE recommends that raised beds, containers, stakes or trellises are made of non-toxic/non-leaching/non-pressure-treated materials. *Corrective action*: replace potentially hazardous materials, or at least use the areas near the potentially hazardous examples for non-food only, such as flowers. Have at least a 1-foot barrier from potential contaminants between potentially hazardous materials and areas where food items are grown.
- 10. Chemicals, including fertilizers, paints, lubricants, cleaning supplies etc. are not stored in close proximity to the garden or to the harvested food. It's best to keep these items in their own specific secured area, such as a locked shed.
- 11. Staff Training: See "School Garden Staff Training & Log"
- 12. Student Training: See "School Garden Student Training & Log"
- 13. Seeds, plants and starts are obtained from reputable sources that are licensed to sell them. Do not accept seeds, plants or food from household sources.
- 14. Consult with the necessary school and district staff. This includes the school principal and the district Integrated Pest Management (IPM) Coordinator (often a facilities manager), and any other staff you are advised to consult with before beginning.
- 15. Pesticides and insecticides are not to be used on school gardens.
 - Other than insecticidal soap, any pesticide products (including over-the-counter baits for ants, rodents, etc.) used on school campuses must be applied by a licensed pesticide applicator and each application must be accompanied by the proper posting and notification as per Oregon's school integrated pest management law (ORS 634.700-634.750). For more information about school IPM, see the Oregon State University School IPM Program page: http://www.ipmnet.org/tim/IPM in Schools/IPM in Schools-Main Page.html

ODE recommends keeping this completed checklist on hand for the current plus three following years.

WEEKLY CHECKLIST (Tab 2):

- Composting: ODE recommends that you do not use cafeteria scraps for composting, UNLESS the garden personnel and monitors are trained in compost management and have a solid system in place that minimizes risks. Raw plant material tends to be very safe, however if the school garden considers bringing in food scraps for the cafeteria the compost will potentially be much more likely to attract pests and are more likely to support the growth of harmful pathogens. A proper composting system includes:
 - a. Training staff and students in proper composting techniques
 - b. Documents time and temperatures required to destroy pathogens associated with composting
 - c. Monitoring of composting stations (see below). Follow the criteria in the "Staff Training" section (Tab 4) for procedures for composting.
- 2. Collecting cafeteria scraps: ensure that the station is monitored (not just left alone in the cafeteria) by an adult, or a trained, responsible student who can identify what should and shouldn't be composted. Typically interest can wane and compost stations can be left unattended. Proper monitoring is crucial to ensure that correct items are composted so that harmful pathogen growth is not supported. Best practices include:
 - a. Have a poster with pictures of items that are acceptable and unacceptable on it.
 - b. During an assembly, have a segment on composting. Students get behind the process and are excited once they know the "whys" and what is going on.
- 3. Outside compost piles attract pests! Best practices for these systems include the closedsystem compost piles that have a lid to reduce pest activity. Buckets that collect and deliver the compost to the pile should be washed/rinsed/sanitized.
- 4. Fertilizers and any soil or amendments treated with chemicals: follow instructions. Only to be applied by adults.
- 5. Is there any evidence of "abuse" from animals? Including sightings of cats using the beds as a litter box, rodent activity, deer activity, etc.
 - a. Corrective action: If there is evidence of animal abuse (feces or urine) in the garden, food in that area should NOT be harvested for human consumption.
 - b. Remove and discard part of plant/soil that is affected by contamination.
 - c. Try to keep animals out by erecting fencing, or traps, etc. There have been success stories with solar-powered electric fencing perimeters to control persistent problems; however you would have to consider that small children are also near these systems, you would need to consider ensuring that precautions were in place to protect the children, like a "caution tape" barrier between kids and the fencing.

- 6. All students/staff have access to restrooms (with potable hot running water/soap/disposable towels.
- Everyone must properly wash their hands with soap and warm water before *harvesting food to be served to the public*. Best practices include having the class line up at the restroom just as they would before lunch service. *NOTE* This process is for HARVESTING food for the cafeteria or public consumption.
- 8. Any students or staff who exhibit symptoms of serious illness (vomiting, fever, or diarrhea) should be prevented from handling produce in the garden so they won't spread harmful bacteria and germs. Typically these individuals shouldn't be at school, but have non-food contact activities for these students and staff in case.
- 9. For safety and sanitation, it is recommended that all students and staff wear close-toed shoes. Flip-flops and bare feet pose physical safety and sanitation issues.
- 10. If manure is used, use only commercially prepared manure on the school garden. This substantially decreases the risk in introducing harmful bacteria to the food.
- 11. Irrigation (for watering) for the school garden should be from a municipal source. If any other source of water is used, including well water or a rain collection system it should be tested (see startup checklist) twice a year. You *SHOULD NOT* use water that's collected from parking lots, gray water, waste water or recycled water from these sources. Water collecting containers should be cleaned and sanitized on a regular basis.
- 12. Insecticides/pesticides should not be used in a school garden. The application of these anywhere on school campuses is regulated by state law. A district's IPM Coordinator must be contacted for pest issues, and situations requiring the use of pesticide products on school grounds.
- 13. Containers used to harvest/transport food should be non-porous, food grade, and easy to sanitize. Do not use: burlap bags; single-use grocery bags; wicker baskets. For assistance with specific examples of items that are safe to use, contact your local food service/cafeteria.
- 14. USDA "Best Practices for Handling Fresh Produce in Schools" **(Tab 3)** should be followed for harvesting/cleaning the produce (attached in this guide). It is very common for the garden staff to harvest the food and deliver it to the cafeteria staff. The cafeteria staff in Oregon is trained in handling the produce correctly (if they are a part of the National School Lunch Program). If they are not available for a harvest, the garden coordinator needs to follow the procedures contained in the handout.
- 15. If the produce will be harvested to be consumed at a later time, the harvested items are to be properly labeled and refrigerated or appropriately stored following the principles in the attachment listed in #13.
- 16. All tools and utensils need to be properly cleaned and free of dirt at the end of the day. Any tools/utensils that come into contact with food need to be properly cleaned and sanitized:
 - a. Scrape any soil/debris off tool
 - b. wash with warm soapy water
 - c. rinse
 - d. Sanitize using proper sanitizer, using iodine, quaternary or chlorine bleach

e. Use corresponding test strip to ensure proper strength of sanitizer. Contact your cafeteria to find out how to purchase test strips and sanitizer, as every cafeteria utilizes them.

Keep these completed weekly logs for the current plus three years.

STAFF TRAINING & LOG (Tab 4):

- It is recommended that at least one person (typically the School Garden Coordinator) attend a school garden certification course. These courses take place in the summer typically, and are usually four or five days in length. They provide overall garden management training and have a garden food safety component.
- All staff members should be trained using the *staff training and log* before working in the garden. Have the staff sign the training log and keep for one year.
- Complete yearly, retain for current plus three years.

STUDENT TRAINING & LOG:

- All students should be trained using the *student training and log* before working in the garden. Have the students sign the training record-or, have the teacher indicate the classroom that was trained, and keep for one year.
- Retain for current plus three years.

COMPOSTING LOG(S):

- Depending on whether you utilize hot or cold composting (or both), it is crucial that you document the proper procedure and process you follow by utilizing these logs.
- Retain for current plus three years

SPECIAL THANKS to the following people in developing of this program:

Caitlin Blethen, Youth Grow Manager, Growing Gardens, Portland OR Chris Bosak, Oregon Department of Education Child Nutrition Services Kari Christensen, Brownfields Program Coordinator, Oregon Health Authority Public Health Division Anne Donahue, City of Eugene Oregon Compost Program Patty Driscoll, Compost Specialist Committee Chair, Oregon State University Extension Service William Fargo, Food Safety Specialist, Food Safety Division, Oregon Department of Agriculture Lisa Carolina Gonzalez, University of Maryland Extension Kristin Hall, Child Nutrition Specialist, Oregon Department of Education Matt Hoffman, MPH REHST, Health Educator, Registered Environmental Health Specialist Trainee, Multnomah County Environmental Health Stewart Jacobsen, Food Safety Projects Coordinator, Agricultural Consultation and Training, Arizona Department of Agriculture Ashley Jeppson, Nevada Department of Agriculture Megan Kemple, Willamette Farm and Food Coalition Gail Langellotto, Ph.D. Statewide Coordinator, Oregon State University Extension Master Gardener Program Andrew Nowak, Slow Food Denver Michelle Markesteyn Ratcliffe, Oregon Department of Agriculture John Moriarty, School Garden Project of Lane County Katy Pelissier, Farm to School Coordinator, Ecotrust Ashley Schimke, Farm to School Specialist, Arizona Department of Agriculture Jennifer Snyder, Faculty Research Assistant, School IPM Specialist, Oregon State University Erica Steinhart, Farm to School Specialist, DC Office of the State Superintendent of Education Anne Wilson, Denver Public Schools Jennifer Young, Nutrition and Physical Activity Coordinator, Oregon Health Authority

Oregon Department of Education School Garden Food Safety Program ~Start-Up Checklist~



REVISED: 10/17/2013 COMPLETE TWICE A YEAR

NO	ITEMS TO CHECK		YES	NO	N/A
1		by See 20)			
	FALL (DATE) (complete	e by Sep. 30)			
2	SPRING (DATE) (complete Identify Garden Coordinator, person who is in charge of this garden	by April 30)			
2	(Name, Title):				
3	At least one designated adult with a current county food handler certifica from the garden served at school, including harvesting, washing, prepari (Name/Title):			ion of f	ood
4	Soil: Identified soil history of the garden site (only needs to be done onc	e)			
5	Soil : Soil has been tested for lead, arsenic and other major metals and to be done once unless reintroducing more soil at a later date) <i>Attach test</i>	contaminants. (needs st results to this form.			
6	Irrigation: is municipal water used?				
7	If not, water is tested for total coliform bacteria at least once per year results to this form.				
8	Growing plot is positioned so that it is not in the path of runoff from agricultural areas, parking lots and roads, or other sources of potential contamination.				
9	Raised beds, containers, stakes or trellises are made of non-toxic/non leaching/non pressure- treated materials. Corrective action: for non-food (flowers) or at least a 1-foot non-food barrier.				
10	Chemicals, including fertilizers, paints, lubricants, cleaning supplies etc. are not stored in close proximity to the garden or to the harvested food				
11	TRAINING : All garden staff have completed Staff Training and Log before working in the garden and supervising students.				
12	TRAINING: All students have completed Student Training and Log before working in the garden.				
13	Seeds and/or plants are procured from reputable sources				
14	You have connected with the district Facilities person / integrated pest m Coordinator to discuss proper location of garden and any compost areas				
Corre	ective Action:				

Oregon Department of Education School Garden Food Safety Program ~Start-Up Checklist~



REVISED: 10/17/2013 COMPLETE TWICE A YEAR

Corrective Action Continued:

Oregon Department of Education School Garden Food Safety Program ~Weekly Garden Checklist~



REVISED: 10/17/2013

NO	ITEMS TO CHECK	YES	NO	N/A
	WEEK OF (DATE)			_
1	Employing Best Composting Practices (Lane County Extension Service Compost Specialist Program –see staff training section/tab 4): Compost log is completed as to instructions on the log			
2	Compost collection station in cafeteria is staffed and monitored by an adult or a trained student			
3	Label instructions for the use of soils and fertilizers are being followed. Fertilizers are only applied by adults.			
4	There is no evidence of abuse from domestic and/or wild animals			
5	All Students/Staff have access to restrooms (with potable hot running water/soap/paper towels)			
6	Proper personal hygiene practices are in place: All students/staff wash hands before harvesting food for public.			
7	Persons who are ill (vomiting, fever, diarrhea) are prevented from working in a garden and handling food			
8	All students/staff should wear closed-toed shoes.			
9	Untreated (not commercially prepared) manure is not being used			
10	Gray water, waste water, recycled water from any source, or runoff water from parking lots is not being used			
11	Insecticides and/or pesticides are not used			
12	Containers used to transport harvested items are food-grade, properly cleaned and in good condition			
13	USDA "Best Practices for Handling Fresh Produce in Schools" (vegetable harvesting/cleaning) procedures are being followed for those items being used in cafeteria or otherwise consumed. (<u>www.nfsmi/producesafety.org</u> – search "handling fresh produce")			
14	Harvested items are labeled and properly stored prior to use in cafeteria or otherwise consumed.			
15	At end of the day, tools are properly cleaned and stored. Tools and utensils that come in contact with food are properly cleaned and sanitized			
	RECTIVE ACTION for any items marked "NO" above: Use other side. See companion g	guide fo	or help	on
	ctive actions. RECTIVE ACTION			
COR	ACTIVE ACTION			

Oregon Department of Education School Garden Food Safety Program ~Weekly Garden Checklist~



REVISED: 10/17/2013

Corrective Action Continued



BEST PRACTICES HANDLING FRESH PRODUCE IN SCHOOLS

Fruits and vegetables are an important part of a healthy diet. Introducing children to them in schools will improve their present and future health. Fresh produce must be handled safely to reduce the risks of foodborne illness. There are a number of steps that foodservice employees can take to minimize the chances for fruits and vegetables they handle to become contaminated. Best practices for handling all types of produce are described in this fact sheet, along with practices specific to leafy greens, tomatoes, melons, and sprouts.

Contamination of produce with harmful microorganisms can occur at all stages of production, processing, transportation, storage, preparation, and service. To prevent foodborne illness, fresh produce needs to be handled with care at each step from farm to table.

Recommendations For Handling Fresh Produce

PURCHASING AND RECEIVING

- Use purchasing specifications that include food safety requirements, such as maintaining produce at the proper temperature, maintaining clean and pest-free storage areas and delivery vehicles, and complying with federal and state food safety laws and regulations.
- Ensure suppliers are getting produce from licensed, reputable sources.
- Check storage and handling practices of vendors.
- Establish procedures for inspecting and accepting or rejecting incoming deliveries. Procedures should include checking the condition of the fresh produce and the transportation vehicles to make sure specifications are met.

WASHING AND PREPARATION

- Inspect produce for obvious signs of soil or damage prior to cutting, slicing, or dicing.
 When in doubt about damaged produce, either cut away the affected areas or do not use the item.
- Wash produce before serving or cutting using either:

Continuous running water.

Chemical disinfectants, used according to the manufacturer's label instructions for recommended concentration and contact time. *Note: Do not soak produce or store in standing water.*

- Do not rewash packaged produce labeled "ready-to-eat," "washed," or "triple washed."
- Wash thoroughly with hot soapy water all equipment, utensils, and food contact surfaces that come into contact with cut produce. Rinse, sanitize, and air-dry before use.

HAND HYGIENE

- Wash hands thoroughly with soap and water before handling or cutting fresh produce. Rewash hands after breaks, visiting restrooms, sneezing, coughing, handling trash or money, or anytime hands become soiled or otherwise contaminated.
- Use a barrier such as gloves, deli paper, or an appropriate utensil to touch ready-to-eat produce. *Note: This does not eliminate the need for frequent proper handwashing.*
- Always wash hands before putting on disposable gloves.
- Change disposable gloves anytime the gloves may have been contaminated or when changing tasks.
- Do not wash or reuse disposable gloves.
- Change disposable gloves if they are torn or damaged.

SERVING

- Do not store produce in direct contact with ice or water while on display on serving lines and salad bars.
- Mark the time when cut produce is displayed without refrigeration. Display cut produce for a maximum of 4 hours if not in a refrigeration unit or containers surrounded by ice. Discard any uneaten produce at the end of 4 hours.
- Create safe salad bars and self-service lines by taking the following actions:
 - Protect food with sneeze guards or food shields in a direct line between the food and the mouth or nose, usually 14 to 18 inches above the food.
 - Use cleaned and sanitized long-handled tongs, spoons, and ladles so bare hands do not touch food and the utensils do not drop into the serving pans.
 - Change utensils periodically.
 - Set up the salad bar or self-service line as close to mealtime as possible to reduce the time that produce sits out.
 - Keep cold foods at or below 41°F in a refrigeration unit or surrounded by ice.
 - Monitor and document the internal temperature of self-service items every 30 minutes as with other foods on the service lines.
 - Clean up spills promptly. Wiping cloths should be stored in sanitizing solution and laundered daily.
 - Teach children salad bar etiquette. Assign an adult to monitor the salad bar and self-service line to make sure the customers—especially children—are not touching food with their hands, tasting food while in line, putting their heads under the sneeze guard, or returning food items.
 - Clearly label all salad dressings and other containers to discourage tasting.
 - Never add freshly prepared food to food already on salad bars and self-service lines.

STORAGE

- Maintain produce at the temperature recommended for the variety and particular stage of ripeness.
- Store produce at least 6 inches off the floor, including in walk-in refrigerators.
- Store produce in a covered container and above other items that might cause contamination.
- Follow manufacturer's instructions for the product such as "keep refrigerated" or "best if used by."
- Establish a policy for produce that is cut in-house to specify how long the refrigerated cut product may be used. Mark the product with "prepared on" or "use by" date.
- Wash produce just before preparation, not before storage.

TRAINING AND GENERAL FOOD SAFETY PRACTICES

- Develop training programs to teach the importance of food safety and proper handling of produce to all food handlers.
- Practice good food safety and food handling techniques to prevent cross-contamination.

Recommendations For Specific Types Of Produce



MELONS

- Avoid using whole melons that have visible signs of decay or damaged rinds (such as mechanical damage or cracking) due to the increased risk that harmful bacteria may have contaminated the melons.
- Wash the outer surface of the melon thoroughly under running cool tap water to remove surface dirt. Scrub melons with a clean produce brush before cutting. Cut away any bruised or damaged areas before serving.
- Discard cut melons after 4 hours if maintained at 41°F or above. If possible, display cut melons in a refrigerated case, not just on top of ice.
- Display cut melons for a maximum of 4 hours without being kept cool with refrigeration or ice and discard uneaten melons at the end of 4 hours.
- Mark the date on refrigerated cut melons to indicate that they must be consumed or discarded within 7 days.

TOMATOES

- Do not wash tomatoes in cold water. Use wash water temperatures that are at least 10°F warmer than the internal tomato temperature to prevent exterior bacteria from entering the interior of the tomato during washing.
- Ensure whole tomatoes are free from obvious signs of soil and skin damage, such as punctures, prior to cutting, slicing, or dicing. Either cut away any bruised or damaged areas, or do not use the tomato.
- Hold tomatoes at 41°F or below after cutting, including during display on serving lines and salad bars.
- Ensure the temperature of tomatoes purchased as fresh-cut (i.e., sliced, diced, or chopped) is 41°F or lower upon delivery and the tomatoes were kept cool continuously during transport. Reject fresh-cut tomatoes delivered at a temperature higher than 41°F.
- Mark the date on refrigerated cut tomatoes to indicate that they must be consumed or discarded within 7 days.
- Do not store cut tomatoes in direct contact with ice or water.



LEAFY GREENS

- Do not use leafy greens with visible signs of decay or damage because there is an increased risk of the presence of harmful bacteria. When in doubt about the use of decayed or damaged product, either remove the unusable portions or do not use the leafy greens.
- Do not rewash packaged produce labeled "ready-to-eat," "washed," or "triple washed."

SPROUTS

Due to the increasing number of illnesses associated with eating raw sprouts, the Food and Drug Administration has advised all consumers—especially children, pregnant women, the elderly, and persons with weakened immune systems—to not eat raw sprouts as a way to reduce the risk of foodborne illness. All sprouts should be cooked thoroughly before eating to reduce the risk of illness.

Resources

Council for Agricultural Science and Technology. **Food Safety and Fresh Produce: An Update.** Available at http://www.cast-science.org/publications.asp

Food and Drug Administration. **Draft Guidance for Tomatoes, Leafy Greens, and Melons.** Available at http://www.fda.gov/Food/FoodSafety/Product-SpecificInformation/FruitsVegetablesJuices/FDAProduceSafetyActivities/ucm174086.htm

Food and Drug Administration. Safe Handling of Raw Produce and Fresh-Squeezed Fruit and Vegetable Juices. Available at http://www.cfsan.fda.gov/~dms/prodsafe.html

> National Restaurant Association. **Guidelines on How to Keep Salad Bars Safe.** Available at http://www.restaurant.org/foodsafety/how_to_salad.cfm

U.S. Department of Agriculture. **Fresh Fruit and Vegetable Program Handbook.** Available at http://www.fns.usda.gov/cnd/FFVP/Resources/FFVPhandbookFINAL.pdf

U.S. Department of Agriculture. **Fruits & Vegetables Galore: Helping Kids Eat More.** Available at http://www.fns.usda.gov/TN/Resources/fv_galore.html

This project has been funded at least in part with funds from the USDA Food and Nutrition Service. The content of this publication does not necessarily reflect the views or policies of the Department, nor does mention of trade names, commercial products, or organizations imply endorsement by the U.S. Government.

The University of Mississippi is an EEO/AA/Title VI/Title IX/Section 504/ADA/ADEA Employer.



Please feel free to reproduce and distribute this publication. Copies are also available on our Web site: www.nfsmi.org Information about this and other topics may be obtained by contacting the NATIONAL FOOD SERVICE MANAGEMENT INSTITUTE, The University of Mississippi; Telephone: 800.321.3054; Item number ET100-10



Oregon Department of Education

School Garden Food Safety Program

Staff Training Manual and Log



- Someone who is in charge of food from the garden that is going to be harvested, prepared and served to the public MUST have a current food handler's card. This person is usually the kitchen manager, but it's recommended that the garden coordinator get one so they know the proper food handling procedures. They are inexpensive and available at <u>www.orfoodhandlers.com</u>.
- ODE recommends that (at least) the Garden Coordinator be certified through a school garden training certification course.
 - In Oregon, *Growing Gardens* in Portland hosts school Garden Coordinator Certificate Trainings. They are typically held in the summer, and are in multiple locations throughout the State. You can find out all information by going to www.growing-gardens.org, and then go to the "resources" tab, and scroll down to "school gardens."
 - Another option is the Oregon 4-H Gardens Teacher and Volunteer Winter Training by Oregon State University in Multnomah County. Contact Maureen Hosty at 503-916.6075. Website: Google "Oregon 4H school garden teacher and volunteer training"
 - Cornell online training / GHP GAP (good handling practices/good agricultural practices <u>www.gaps.cornell.edu</u> -\$190
- The Garden Coordinator, Kitchen Manager or Cook, Food Service Director, Principal, and any other key stakeholders (like Facilities folks / Integrated Pest Management (IPM) Coordinator) should meet at the beginning of the school year so they all have a common understanding and agree about all things relating to the garden. The items discussed should be things such as: who is responsible for cleaning harvest containers; compost station locations; delivery times of produce; etc.
- The following are general guidelines that should be followed to ensure food safety in your garden. Remember, children are in an age group that is highly susceptible to foodborne illness.

When a school serves food from a school garden to students or the public, there are the same concerns as a cafeteria purchasing from a distributor or farm. There is an opportunity to make people sick if simple procedures aren't followed.

Hygiene: All students/staff wash hands thoroughly with soap and water before harvesting food to be served to the public. When harvesting, rewash hands after breaks, visiting restrooms, sneezing, coughing, handling trash or money, or anytime hands become soiled or otherwise contaminated. After produce is washed and ready to be served in the cafeteria, barriers such as gloves, deli paper or appropriate utensil should be used to touch ready-to-eat produce.

SPECIFICALLY, here is the procedure for hand washing when serving food to the public:

- Wash hands with warm, soapy water for 20 seconds
- Before and after handling food
- After using the bathroom*
- After blowing your nose, coughing or sneezing*
- Dry hands with single-use, disposable towels

*Note: when handling ready to eat food, the Oregon food handling rules state that after the above actions occur (bathroom break/blowing nose) the food handler must double-wash their hands. This means washing their hands in the restroom, and again at the hand washing station when returning to the worksite, typically a cafeteria. However, the Oregon health codes also stipulate that some workplaces will probably NOT have a proper hand washing sink available – like in a garden. In this case, the second step is not required, however harvester must take special attention and make sure they do an adequate job after going to the restroom or blowing their nose.

It's crucial that ANY student or staff be excluded from gardening activities if they are ill, exhibiting symptoms such as vomiting, diarrhea or fever, so they will not spread bacteria leading to foodborne illness. Typically, students who are ill should not be at school anyway, so this shouldn't be an issue.

Composting: Composting cafeteria scraps can bring serious problems to the school. Compost piles can attract pests and potentially introduce harmful bacteria if the system is not done properly and monitored very carefully. **ODE recommends that you do not use cafeteria scraps for composting**, <u>UNLESS</u> your school garden staff (and students involved) are properly trained and have a solid system in place to ensure that you are minimizing risks associated with composting.

- One effective form of a cafeteria compost station is a table with holes cut into them and 5-gallon buckets placed in the holes.
- Have a poster with pictures available showing what to compost and NOT to compost. The provided examples are available on our website:
 <u>www.ode.state.or.us/go/f2sgardens</u>. Click on the "School Garden" box and scroll down to the "Garden Safety" section. School assemblies work very well to educate the

students all at once, showing examples of what's ok to compost. Composting can generate excitement about the garden, too.



- Have a trained adult or *responsible* student monitoring the composting station during meal service period. Responsible students should be trained on this composting system.
- After the material is delivered to the compost pile, the buckets need to be cleaned and sanitized.
- <u>Do not compost</u>: bread & rice (doesn't break down in the same way as others, and introduces harmful molds); meat products; dairy products; heavily coated or printed paper; magazines; oil of any kind; animal feces; sawdust (unless you know if all of the wood was untreated); diseased plants; weeds that are spread by roots or gone to seed.
- <u>OK to compost</u>: fruit and vegetable food scraps that are free of dressing or oils; banana peels; orange peels; melon rinds; apple cores; napkins; grass clippings; leaves; straw; shredded cardboard; coffee grounds and coffee filters.
- Although most all materials will eventually decompose, an efficient compost pile should have a 2:1 ratio of "browns" (leaves, cardboard, etc.) to "greens" (grass clippings, veggies) (known as a "C:N" ratio carbon/nitrogen).
- Compost procedure:
 - Complete the composting log (included in this guide), following these procedures:
 - For "hot composting," internal temperature should read at least 131 degrees for three days, then turned two times and reach 131 degrees after each turn at least.
 - Do not let the pile get above 140 degrees
 - The pile should then be cured for 3 months.
 - Note: The pile needs to be three cubic feet in size for doing hot compost and you will need to adjust the C:N ratio to reach the required temperatures to kill pathogens. Consult your OSU extension office for more guidance.

- Composting thermometer used should be regularly calibrated. To calibrate a bimetallic stem thermometer, fill a cup with crushed ice, and then top it off with water. Immerse the end of the thermometer in the cup and stir around. The needle should read at exactly 32 degrees. Adjust the nut on the underside of the dial until this number is reached. Document when you do this procedure on the composting log.
- For "cold" composting, pile should be protected from pests and cured for at least one year after the last time last is added to the pile.
- Cold composting can take up to two years to completely break down!

Harvesting: Containers must be easily cleanable, non-porous, and food-grade. All implements and containers that come in contact with produce must be washed/sanitized.

Preparing harvested fruits and veggies: Typically harvested food from the garden is delivered to the cafeteria via a prior arrangement. The cafeteria staff would then prepare the produce, but in many cases this might be done by the garden staff. One of the most common problems is that school garden produce typically has a lot more dirt and bugs associated with the produce than produce purchased from conventional sources. This is why it's crucial to pay special care in cleaning and prepping. Produce must be washed THOROUGHLY! Cafeteria staff sometimes has to double/triple wash school garden produce.

Animal activity: Including rodents, cats, and deer. It's important to remember that a typical garden is in the animal's natural environment. It's almost impossible to keep animals completely out of the garden; however fencing or cages should reduce animal activity. Animals can sometimes jump over the tallest fencing. Other considerations to keep in mind:

- You might want to make a small-gauge chicken wire or similar fencing to keep rodents from getting through easily.
- Also, burying the wire a few inches into the ground discourages animals to burrow under.
- Netting to discourage deer and birds
- It is important to realize that you cannot totally keep out all pests from a garden no matter what steps you take! It may be possible to minimize the pest activity by following some of the steps listed above. Consider "flagging" areas where animal abuse occurs, remove plant and dirt that's been contaminated.

Eating "right out of the garden:" Only produce that is not in direct ground contact and shows no obvious signs of soil or other contamination should be eaten directly ("grazing"). Only the Garden Coordinator or responsible adult in charge should tell kids exactly what is safe to eat.

There could potentially be something that would be unsafe to eat in a garden, such as rhubarb leaves.

Tools: Any use of tools (shovels, rakes, clippers, etc.) should be closely and directly supervised by adult(s) in charge. Power equipment such as tillers or trimmers are not allowed to be used by students under the age of 18 by Oregon law.

TRAINING DATE:	Person who conducted tra	aining:	
The following people attended a training session on School Garden Food Safety:			

Attach as many sheets as necessary if more are needed. Retain for current plus three following years.

Oregon Department of Education

School Garden Food Safety Program

Student Training Manual and Log



- Identify the Garden Coordinator. This person is in charge of the garden. All students MUST follow their direction in order to keep students safe.
- **Kitchen Manager or Cook**. Identify that person if the kids are dropping food off in the kitchen. The same goes for this person: They are in charge of the kitchen and all students must follow their direction in order to keep students and food safe.

All students MUST properly **wash their hands** before harvesting produce to be served to the public from the garden. This is to be done thoroughly with soap and warm water (*See below). Rewash hands after breaks, visiting restrooms, sneezing, coughing, handling trash or money, or anytime hands become otherwise contaminated. After produce is washed and ready to be served in the cafeteria or garden, barriers such as gloves, deli paper or appropriate utensil should be used to touch ready-to-eat produce. *SPECIFICALLY, here is the procedure for hand washing when serving food to the public in the garden*:

- Wash hands with warm, soapy water for 20 seconds
- Before handling food
- After using the bathroom*
- After blowing your nose, coughing or sneezing*
 - *See note on "double hand washing" in Staff Training section
- Dry hands with single-use, disposable towels
 - *NOTE: in most cases, there won't be warm water available in the garden. Discretion will need to happen on the part of the garden coordinator. A handwashing station or simply a garden hose will be better than nothing if hands become soiled.
- Tell students that when garden produce is harvested, the food going to the cafeteria is the same as any food going to a restaurant. They wouldn't want people to touch their hair or face before they touched their food, right? In the garden, students shouldn't touch their face/hair and then touch food that is going to the cafeteria.
- **Shoes**: Flip-flops or bare feet can be hazardous in a garden, as well as unsanitary. Closed-toed shoes should be worn in the garden.
- Eating food right from the garden. NOTHING should be eaten without asking the Garden Coordinator!
- **Tools:** The adult in charge will supervise any use of tools. Only use the tools in their intended use.
- Fertilizers: Only adults are allowed to handle and apply fertilizers.
- Insecticides/Herbicides: No insecticides or herbicides should be used.

• **Composting**. Some of the older students may be asked to assist with the compost station in the cafeteria. They will have to be trained to know what is safe to compost and what is NOT allowed in the compost bucket. It's suggested that the students follow the training in the "staff training" section. Students who are responsible to monitor the station need to pay close attention to the station, as it can be a very busy time.

TRAINING DATE: Person who conducted training:			
The following students or classrooms have been through the School Garden Food Safety			
Training:			

Attach as many sheets as necessary if more are needed. Retain for current plus three following years.

Compost log: Hot Composting

Keep on file for current plus three years. Instructions:

- Internal temperature should be read at least 131 degrees for three days, then turned two times and reach 131 degrees after each turn at least.
- Do not let the pile get above 140 degrees
- The pile should then be cured for 3 months
- Note: The pile needs to be three cubic feet in size for doing hot compost and you will need to adjust the C:N ratio to reach the required temperatures to kill pathogens, and minimize flies and other pests. Consult your OSU extension office for more guidance.
- Composting thermometer used should be regularly calibrated. Use this form for documenting when you calibrate the thermometer. See staff training guide for details.

Action	Date	Temperature

Action	Date	Temperature

Retain for current plus three following years

Compost log: Cold Composting

Keep on file for current plus three years.

Instructions:

- Pile should be protected from pests and cured for at least one year after the last time last is added to the pile.
- Cold composting can take up to two years to completely break down.

Action	Date	Temperature

Retain for current plus three following years